arXiv:quant-ph/0507227v2 4 Jan 2006

Violating Bell Inequalities Maximally for Two d-Dimensional Systems

Jing-Ling Chen,[1,2,*] Chunfeng Wu,[2] L.C. Kwek,[2,3] C.H. Oh,[2,†] and Mo-Lin Ge[1]

[1] *Theoretical Physics Division, Chern Institute of Mathematics, Nankai University, Tianjin 300071, P.R.China*
[2] *Department of Physics, National University of Singapore, 2 Science Drive 3, Singapore 117542*
[3] *Nanyang Technological University, National Institute of Education, 1, Nanyang Walk, Singapore 637616*

(Dated: February 1, 2008)

We investigate the maximal violation of Bell inequalities for two d-dimensional systems by using the method of Bell operator. The maximal violation corresponds to the maximal eigenvalue of the Bell operator matrix. The eigenvectors corresponding to these eigenvalues are described by asymmetric entangled states. We estimate the maximum value of the eigenvalue for large dimension. A family of elegant entangled states $|\Psi\rangle_{\rm app}$ that violate Bell inequality more strongly than the maximally entangled state but are somewhat close to these eigenvectors is presented. These approximate states can potentially be useful for quantum cryptography as well as many other important fields of quantum information.

PACS numbers: 03.65.Ud, 03.67.-a,42.50.-p

The famous Clauser-Horne-Shimony-Holt (CHSH) inequality [1, 2] for two entangled spin-1/2 particles has always provided an excellent test-bed for experimental verification of quantum mechanics against the predictions of local realism [3]. It is well-known that all pure entangled states of 2 dimension violate the CHSH inequality: the maximum quantum violation of $2\sqrt{2}$ being often called the Tsirelson's bound [4]. In 2000, contrary to previous study, Kaszlikowski *et al* showed numerically, based on linear optimization techniques, that the violations of local realism increase with dimensions for two maximally entangled d-dimensional systems (qudits) ($3 \leq d \leq 9$) [5]. A year later, Durt *et al* extended the analysis to $d = 16$ under special experimental settings[6]. In the same year, an analytical proof was constructed for two maximally entangled 3-dimensional systems (qutrits) [7]. In 2002, two research teams independently developed Bell inequalities for high-dimensional systems: the first one is a Clauser-Horne type (probability) inequality for two qutrits [8]; and the second one is a CHSH type (correlation) inequality to two arbitrary d-dimensional systems [9], now known as the Collins-Gisin-Linden-Masser-Popoescu (CGLMP) inequalities. The tightness of the CGLMP inequality was demonstrated in Ref. [10]. The maximally entangled state of two-qudit reads $|\Psi\rangle_{\rm mes} = \frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} |jj\rangle$ where $|j\rangle$ is the orthonormal base in each subsystem. Collins *et al* restricted their investigation to $|\Psi\rangle_{\rm mes}$, because on one hand $|\Psi\rangle_{\rm mes}$ is a simple state, on the other hand there is a very natural viewpiont that maximally entangled states would maximally violate the Bell inequality, just as the CHSH inequality has worked for two-qubit. The results of [9] was numerically consistent with Ref. [5]. For $|\Psi\rangle_{\rm mes}$, the Tsirelson's bound or Bell expression I_d is given by [9]

$$\begin{aligned} I_d(|\Psi\rangle_{\rm mes}) &= 4d \sum_{k=0}^{\ell} (1 - \frac{2k}{d-1}) (\frac{1}{2d^3 \sin^2[\pi(k+\frac{1}{4})/d]} \\ &\quad - \frac{1}{2d^3 \sin^2[\pi(-k-1+\frac{1}{4})/d]}), \end{aligned} \tag{1}$$

showing that the maximum quantum violation increases with the dimension d (here $\ell = [d/2-1]$ represents the integer part of $(d/2-1)$.). When $d \to \infty$, it is interesting to note that

$$\begin{aligned} \lim_{d\to\infty} I_d(|\Psi\rangle_{\rm mes}) &= \frac{2}{\pi^2} \sum_{k=0}^{\infty} [\frac{1}{(k+1/4)^2} - \frac{1}{(k+3/4)^2}] \\ &\simeq 2.96981, \end{aligned} \tag{2}$$

a number related to the Catalan's constant.

Contrary to prevalent belief, Acín *et al* studied the quantum nonlocality of two-qutrit as well as two d-dimensional systems up to $d = 8$ and discovered another unexpected result: there existed non-maximally entangled states that lead to greater violation of the CGLMP inequalities compared with maximally entangled states[11]. This surprising result still lacks of a good intuitive explanation. From the TABLE I of Ref. [11], one observes that the maximal violation increases with dimension d, and it reaches 3.1013 for $d = 8$. This gives rise to a natural question:" What are the maximal violations of the CGLMP inequalities if one increases the dimension d, especially when d goes to infinity?" The purpose of this Letter is to investigate the maximal violations of the CGLMP inequalities for higher dimensional

systems. We also present a family of elegant entangled states $|\Psi\rangle_{\rm app}$ whose corresponding maximal violations are approximate to the real ones.

Before computing the quantum violation, let us first estimate the upper bound under quantum mechanics. Significantly, Ref. [13] has recast the CGLMP inequality into a form that is very similar to the CHSH inequality:

$$I_d = Q_{11} + Q_{12} - Q_{21} + Q_{22} \leq 2, \tag{3}$$

where Q_{ij} are the correlation functions defined by probabilities in the following way

$$Q_{ij} = \frac{1}{S}\sum_{m=0}^{d-1}\sum_{n=0}^{d-1} f^{ij}(m,n)P(A_i = m, B_j = n), \tag{4}$$

where $S = (d-1)/2$ is the spin of the particle for the d-dimensional system, $f^{ij}(m,n) = S - M(\varepsilon(i-j)(m+n), d)$; $\varepsilon(x) = 1$ and -1 for $x \geq 0$ and $x < 0$ respectively; $M(x,d) = (x \ {\rm mod} d)$ and $0 \leq M(x,d) \leq d-1$. Refs. [9] and [13] have proved that $I_d \leq 2$ for hidden variable theory. From Eq. (4), one notes that the extreme values of Q_{ij} are ± 1 for both local realistic description and quantum mechanics, therefore it is impossible that the maximum quantum violation of I_d is larger than 4. Furthermore I_d cannot reach 4, because Q_{ij} are constrained to each other, if three of them are set to be 1, the fourth must also be 1. Consequently, one can conclude easily from above analysis that the maximal quantum violation of I_d is a number between 2 and 4.

To generalize Acín's approach, we first note that it has been shown that unbiased multiport beam splitter [12] can be used to test violation of local realism of two maximally entangled qudits. Unbiased d-port beam splitter is a device with the following property: if a photon enters any of the d single input ports, its chances of exit are equally split among the d output ports. In fact one can always build the device with the distinguishing trait that the elements of its unitary transition matrix T are solely powers of the root of unity $\gamma = \exp(i2\pi/d)$, namely $T_{kl} = \frac{1}{\sqrt{d}}\gamma^{kl}$. In front of i-th input port of the device a phase shifter is placed to change the phase of the incoming photon by $\phi(i)$. These d phase shifts, denoted for convenience as a "vector" of phase shifts $\hat{\phi} = (\phi(0), \phi(1), ..., \phi(d-1))$, are macroscopic local parameters that can be changed by the observer. Therefore, unbiased d-port beam splitter together with the d phase shifters perform the unitary transformation $U(\hat{\phi})$ with the entries $U_{kl} = T_{kl}\exp(i\phi(l))$. Devices (Bell multiports) endowed with such a matrix were proposed, and readers who are interested in it can refer to Refs. [5][12]. The approach developed in [11] is related to Bell operator. An arbitrary entangled state of two-qudit reads

$$|\Phi\rangle = \sum_{j,j'=0}^{d-1} \alpha_{jj'}|jj'\rangle. \tag{5}$$

The quantum prediction of the joint probability $P(A_a = k, B_b = l)$ when A_a and B_b are measured in the initial state $|\Phi\rangle$ is given by

$$\begin{aligned} P(A_a = k, B_b = l) &= {\rm Tr}[(U(\hat{\phi}_a)^\dagger \otimes U(\hat{\varphi}_b)^\dagger)\ \hat{\Pi}_k \otimes \hat{\Pi}_l\ (U(\hat{\phi}_a) \otimes U(\hat{\varphi}_b))\ |\Phi\rangle\langle\Phi|] \\ &= \frac{1}{d^2}\sum_{j,j',m,m'=0}^{d-1} \alpha_{jj'}\alpha^*_{mm'} e^{i[\phi_a(j)+\varphi_b(j')+\frac{2\pi}{d}(jk-j'l)-\phi_a(m)-\varphi_b(m')-\frac{2\pi}{d}(mk-m'l)]}, \end{aligned} \tag{6}$$

where $\hat{\Pi}_k = |k\rangle\langle k|$, $\hat{\Pi}_l = |l\rangle\langle l|$ are the projectors for systems A and B, respectively. Substituting Eq. (6) into the CGLMP inequality, one gets the Bell expression for the state $|\Phi\rangle$:

$$
\begin{aligned}
I_d(|\Phi\rangle) = \quad & \frac{1}{d^2}\sum_{j,j',m,m'=0}^{d-1}\alpha_{jj'}\alpha^*_{mm'}\sum_{l=0}^{d-1}e^{i\frac{2\pi}{d}[(j-m)-(j'-m')]l} \\
& \times\{e^{i[\phi_1(j)-\phi_1(m)+\varphi_1(j')-\varphi_1(m')]}\sum_{k=0}^{\ell}(1-\frac{2k}{d-1})(e^{i\frac{2\pi}{d}k(j-m)}-e^{-i\frac{2\pi}{d}(k+1)(j'-m')})+ \\
& e^{i[\phi_1(j)-\phi_1(m)+\varphi_2(j')-\varphi_2(m')]}\sum_{k=0}^{\ell}(1-\frac{2k}{d-1})(e^{-i\frac{2\pi}{d}k(j'-m')}-e^{i\frac{2\pi}{d}(k+1)(j-m)})+ \\
& e^{i[\phi_2(j)-\phi_2(m)+\varphi_1(j')-\varphi_1(m')]}\sum_{k=0}^{\ell}(1-\frac{2k}{d-1})(e^{-i\frac{2\pi}{d}(k+1)(j'-m')}-e^{i\frac{2\pi}{d}k(j-m)})+ \\
& e^{i[\phi_2(j)-\phi_2(m)+\varphi_2(j')-\varphi_2(m')]}\sum_{k=0}^{\ell}(1-\frac{2k}{d-1})(e^{i\frac{2\pi}{d}k(j-m)}-e^{-i\frac{2\pi}{d}(k+1)(j'-m')})\}. \qquad (7)
\end{aligned}
$$

$I_d(|\Phi\rangle)$ can be expressed as

$$
I_d(|\Phi\rangle) = \mathrm{Tr}(\hat{B}|\Phi\rangle\langle\Phi|) = \langle\Phi|\hat{B}|\Phi\rangle, \qquad (8)
$$

where $\hat{B}$ is the so-called Bell operator. Starting with the CGLMP inequality and choosing suitable the experimental settings [6][11]

$$
\phi_1(j) = 0, \quad \phi_2(j) = \frac{j\pi}{d}, \quad \varphi_1(j) = \frac{j\pi}{2d}, \quad \varphi_2(j) = -\frac{j\pi}{2d}, \qquad (9)
$$

optimal for maximal violations, we can derive the element of the Bell operator matrix as

$$
\begin{aligned}
B_{mm',jj'} = \quad & \frac{1}{d^2}\sum_{l=0}^{d-1}e^{i\frac{2\pi}{d}[(j-m)-(j'-m')]l} \\
& \{e^{\frac{i\pi}{2d}(j'-m')}\sum_{k=0}^{\ell}(1-\frac{2k}{d-1})(e^{i\frac{2\pi}{d}k(j-m)}-e^{-i\frac{2\pi}{d}(k+1)(j'-m')})+ \\
& e^{-\frac{i\pi}{2d}(j'-m')}\sum_{k=0}^{\ell}(1-\frac{2k}{d-1})(e^{-i\frac{2\pi}{d}k(j'-m')}-e^{i\frac{2\pi}{d}(k+1)(j-m)})+ \\
& e^{\frac{i\pi}{d}(j-m)+\frac{i\pi}{2d}(j'-m')}\sum_{k=0}^{\ell}(1-\frac{2k}{d-1})(e^{-i\frac{2\pi}{d}(k+1)(j'-m')}-e^{i\frac{2\pi}{d}k(j-m)})+ \\
& e^{\frac{i\pi}{d}(j-m)-\frac{i\pi}{2d}(j'-m')}\sum_{k=0}^{\ell}(1-\frac{2k}{d-1})(e^{i\frac{2\pi}{d}k(j-m)}-e^{-i\frac{2\pi}{d}(k+1)(j'-m')})\}, \qquad (10)
\end{aligned}
$$

where $\hat{B}$, in general, is a $d^2\times d^2$ matrix. Ref. [11] found that the maximal eigenvalue of matrix $\hat{B}$ is nothing but the highest quantum prediction of the Bell expression $I_d(|\Phi\rangle_{\mathrm{eig}})$ and the corresponding eigenvector $|\Phi\rangle_{\mathrm{eig}}$ is the state that maximally violates the Bell inequality. Thus, the problem of computing the maximal violation of $I_d(|\Phi\rangle)$ reduces to the determination of the maximal eigenvalue of matrix $\hat{B}$. Due to $\sum_{l=0}^{d-1}e^{i\frac{2\pi}{d}(p-q)l} = d\delta_{pq}$, where $\delta_{pq}=1$ when $p=q$ modulo d and 0 otherwise, the matrix $\hat{B}$ can be further simplified, i.e., it can be decomposed into the sum of d decoupled operators that act individually within the subspaces spanned by the vectors $\{|00\rangle,|11\rangle,...,|(d-1)(d-1)\rangle\}$, $\{|01\rangle,|12\rangle,...,|(d-1)0\rangle\}$,...,$\{|0(d-1)\rangle,|10\rangle,...,|(d-1)(d-2)\rangle\}$, respectively. For the first subspace spanned by the

vectors $\{|00\rangle, |11\rangle, ..., |(d-1)(d-1)\rangle\}$, and with the constraint $j-m = j'-m'$, one can reduce $B_{mm',jj'}$ to

$$\begin{aligned} & B^{red1}_{mj} \\ = & \frac{8}{d}\sin\left[\frac{\pi}{2d}(j-m)\right] \\ & \times\sum_{k=0}^{\ell}(1-\frac{2k}{d-1})\sin\left[\frac{2\pi}{d}(k+\frac{1}{2})(j-m)\right] \\ = & \frac{2}{d-1}\,\frac{1}{\cos[\frac{\pi}{2d}(j-m)]}(1-\delta_{mj}), \end{aligned} \tag{11}$$

where B^{red1}_{mj} is the matrix element located at the m-th row and the j-th column of the reduced Bell operator B^{red1}. Let us denote $B_r = B_{m,m+r}$, since $B_{m,j}$ depends only on the difference $(j-m)$. Therefore, we arrive at the following $d\times d$ real symmetric matrix for the reduced Bell operator:

$$\hat{B}^{red1} = \begin{pmatrix} B_0 & B_1 & B_2 & \dots & B_{d-3} & B_{d-2} & B_{d-1} \\ B_1 & B_0 & B_1 & B_2 & \dots & B_{d-3} & B_{d-2} \\ B_2 & B_1 & B_0 & \ddots & \ddots & \vdots & B_{d-3} \\ \vdots & B_2 & \ddots & \ddots & \ddots & B_2 & \vdots \\ B_{d-3} & \vdots & \ddots & \ddots & B_0 & B_1 & B_2 \\ B_{d-2} & B_{d-3} & \dots & B_2 & B_1 & B_0 & B_1 \\ B_{d-1} & B_{d-2} & B_{d-3} & \dots & B_2 & B_1 & B_0 \end{pmatrix}. \tag{12}$$

One may read from Eq.(11) that B_r are all non-negative numbers, $B_0 \equiv 0$, and interestingly $\lim_{d\to\infty} B_{d-1} = 4/\pi$, $\lim_{d\to\infty} B_{d-j}/B_{d-1} = 1/j$, for $j << d$. The Bell operator $\hat{B}^{red1}$ acts on a matrix column $(a_0, a_1, \cdots, a_{d-1})^T$, which corresponds to the state $|\Psi\rangle = \sum_{j=0}^{d-1} a_j|jj\rangle$ in the first subspace. Actually, an arbitrary state $|\Phi\rangle = \sum_{j,j'=0}^{d-1}\alpha_{jj'}|jj'\rangle$ can always be transformed into its Schmidt decomposition form $|\Psi\rangle = \sum_{j=0}^{d-1} a_j|jj\rangle$ through local unitary transformations, thus it is sufficient to study the maximal violation problem in the first subspace. For a general state $|\Psi\rangle$, one can obtain the Bell expression as

$$\begin{aligned} I_d(|\Psi\rangle) &= \mathrm{Tr}(\hat{B}^{red1}|\Psi\rangle\langle\Psi|) = \sum_{m=0}^{d-1}\sum_{j=0}^{d-1} B_{mj}a_m a_j \\ &= \sum_{r=1}^{d-1} B_r\left(2\sum_{m=0}^{d-1-r} a_m a_{m+r}\right). \end{aligned} \tag{13}$$

For instance, for the maximally entangled state $|\Psi\rangle_{\rm mes}$ with $a_j = 1/\sqrt{d}$, the summation $(\sum_{m,j=0}^{d-1} B_{mj})/d = (\sum_{r=1}^{d-1} 2(d-r)B_r)/d$ recovers the results of Eq. (1).

As the analysis shown in [11], the maximal violation of the CGLMP inequality with the experimental settings (9) corresponds to the maximal eigenvalue of $\hat{B}^{red1}$, and indeed its corresponding eigenvector is a nonmaximally entangled state of two-qudit. For instance, for $d = 3$, one has $B_0 = 0, B_1 = 2\sqrt{3}/3, B_2 = 2$, the eigenvector $|\Psi\rangle^{d=3}_{\rm eig} = \sqrt{\frac{2}{11-\sqrt{33}}}(|00\rangle + \frac{\sqrt{11}-\sqrt{3}}{2}|11\rangle + |22\rangle)$ corresponds to a maximal violation $I_{d=3}(|\Psi\rangle_{\rm eig}) = 1+\sqrt{11/3} \simeq 2.9149$, which is larger than $I_{d=3}(|\Psi\rangle_{\rm mes}) \simeq 2.8729$; for $d = 4$, one has $B_0 = 0, B_1 = 2\sqrt{4-2\sqrt{2}}/3, B_2 = 2\sqrt{2}/3$, $B_1 = 2\sqrt{4+2\sqrt{2}}/3$,the eigenvector $|\Psi\rangle^{d=4}_{\rm eig} = \frac{1}{\sqrt{2+2a^2}}(|00\rangle + a|11\rangle + a|22\rangle + |33\rangle)$, with $a = (\sqrt{2+\sqrt{2}} + \sqrt{8-3\sqrt{2}+4\sqrt{2-\sqrt{2}}} - \sqrt{4+2\sqrt{2}})/(\sqrt{2}+\sqrt{4-2\sqrt{2}}) \simeq 0.73937$, corresponds to a maximal violation $I_{d=4}(|\Psi\rangle_{\rm eig}) = \frac{2}{3}\sqrt{2+\sqrt{2}} + \frac{2}{3}\sqrt{8-3\sqrt{2}+4\sqrt{2-\sqrt{2}}} \simeq 2.9727$, which is larger than $I_{d=4}(|\Psi\rangle_{\rm mes}) \simeq 2.8962$.

By diagonalizing exactly the matrix $\hat{B}^{red1}$, we have obtained the real maximal violations $I_d(|\Psi\rangle_{\rm eig})$ for two entangled qudits. The highest dimension that we have calculated is $d = 8000$. In Fig. 1, one may observe that $I_d(|\Psi\rangle_{\rm eig})$



FIG. 1: Variations of $I_d(|\Psi\rangle)$ with increasing dimension d ($2 \leq d \leq 8000$).

increases with dimension d. Based on the data of $I_d(|\Psi\rangle_{\rm eig})$ from $d = 2$ to $d = 8000$, one has an empirical formula fitting $I_d(|\Psi\rangle_{\rm eig})$ numerically to the dimension d:

$$I_d^{\rm rough}(|\Psi\rangle_{\rm eig}) \ \simeq \ 3.9132 - 1.2891x^{-0.2226} \tag{14}$$

from which one can see that $I_d^{\rm rough}(|\Psi\rangle_{\rm eig}) \simeq 3.9132$ is a coarse-grained limit of the maximal violation for the CGLMP inequality when d tends to infinity.

Analysis of the eigenvectors $|\Psi\rangle_{\rm eig}$ shows that these eigenvectors numerically satisfy some general properties: for instance, $|\Psi\rangle_{\rm eig} = \sum_{j=0}^{d-1} a_j^{\rm eig}|jj\rangle$ with maximal eigenvalue has the following symmetric properties for the coefficients: $a_j = a_{d-1-j}$; and $a_0 : a_1 : a_2 : a_3 : \cdots \simeq 1 : \frac{1}{\sqrt{2}} : \frac{1}{\sqrt{3}} : \frac{1}{\sqrt{4}} : \cdots$ for large d. Thus, we may approximate a family of

elegant entangled states

$$|\Psi\rangle_{\rm app} = \sum_{j=0}^{d-1} a_j^{\rm app}|jj\rangle, \;\; a_j^{\rm app} = \frac{1}{\sqrt{\mathcal{N}}}\,\frac{1}{\sqrt{(j+1)(d-j)}},$$
$$\mathcal{N} = \sum_{j=0}^{d-1}\frac{1}{(j+1)(d-j)} \tag{15}$$

whose corresponding Bell expressions $I_d(|\Psi\rangle_{\rm app})$ are closed to the actual ones $I_d(|\Psi\rangle_{\rm eig})$. For example, for $d = 8000$, the error rate between $I_d(|\Psi\rangle_{\rm eig})$ and $I_d(|\Psi\rangle_{\rm app})$ is only about 0.745%. We have also listed $I_d(|\Psi\rangle_{\rm app})$ and $I_d(|\Psi\rangle_{\rm mes})$ in Table I and drawn the corresponding curves in Fig. 1. One may observe that for $d = 50000$, $I_d(|\Psi\rangle_{\rm mes}) \simeq 2.96981$ has almost reached the limit as shown in Eq. (2); and for $d = 600000$, $I_d(|\Psi\rangle_{\rm app})$ has exceeded 3.80.

In summary, we have investigated the maximal violation of the CGLMP inequalities for two entangled qudits. The maximal violation occurs at the non-maximally entangled state, which is the eigenvector of Bell operator with the maximal eigenvalue. The maximal violations increase with growing dimension d, coming close to 4 when d approaches to infinity. Experimental test has been performed to verify the CGLMP inequalities for the first few high-dimensional systems, and indeed there exist non-maximally entangled states that violate these inequalities more strongly than the maximal entangled ones [14]. Bell inequalities are applicable to quantum cryptography and quantum communication complexity [15], previous researches are mostly based on maximally entangled states. For stronger violation, it may be useful to generate the approximate state in this paper. Nevertheless, it may be significant and interesting to apply the elegant symmetric entangled states $|\Psi\rangle_{\rm app}$ to quantum cryptography as well as other important fields of quantum information.

This work is supported by NUS academic research Grant No. WBS: R-144-000-123-112.

* Electronic address: chenjl@nankai.edu.cn
† Electronic address: phyohch@nus.edu.sg

[1] J. S. Bell, Physics (Long Island City, N.Y.) **1**, 195 (1964).
[2] J. F. Clause, M. A. Horne, A. Shimony and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
[3] A. Aspect, P. Grangier and G. Roger, Phys. Rev. Lett. **47**, 460 (1981); A. Fine, *ibid* **48**, 291 (1982); W. Tittel *et al.*, *ibid* **81**, 3563 (1998); G. Weihs *et al.*, *ibid* **81**, 5039 (1998); M. Rowe *et al.*, Nature **409**, 791 (2001).
[4] N. Gisin, Phys. Lett. A **154**, 201 (1991); A. Peres, Phys. Rev. A **46**, 4413 (1992); N. Gisin and A. Peres, Phys. Lett. A **162**, 15 (1992).
[5] D. Kaszlikowski, P. Gnaciński, M. Żukowski, W. Miklaszewski, and A. Zeilinger, Phys. Rev. Lett. **85**, 4418 (2000).
[6] T. Durt, D. Kaszlikowski, and M. Żukowski, Phys. Rev. A **64**, 024101 (2001).
[7] J.L. Chen, D. Kaszlikowski, L.C. Kwek, C.H. Oh, and M. Żukowski, Phys. Rev. A 64, 052109 (2001).
[8] D. Kaszlikowski, L.C. Kwek, J.L. Chen, M. Żukowski, and C.H. Oh, Phys. Rev. A 65, 032118 (2002); J.L. Chen, D. Kaszlikowski, L.C. Kwek, and C.H. Oh, Mod. Phys. Lett. A **17**, 2231 (2002).
[9] D. Collins, N. Gisin, N. Linden, S. Massar, and S. Popescu, Phys. Rev. Lett. **88**, 040404 (2002).
[10] Lluis Masanes, Quantum Information and Computation, Vol.3, No.4, 345 (2002)
[11] A. Acín, T. Durt, N. Gisin, and J. I. Latorre, Phys. Rev. A **65**, 052325 (2002).
[12] M. Żukowski, A. Zeilinger, and M. A. Horne, Phys. Rev. A **55**, 2564 (1997).
[13] L. B. Fu, Phys. Rev. Lett. **92**, 130404 (2004).
[14] A. Vaziri, G. Weihs, and A. Zeilinger, Phys. Rev. Lett. 89, 240401 (2002).
[15] A. K. Ekert, Phys. Rev. Lett. 67, 661 (1991); D. Kaszlikowski, D. K. L. Oi, M. Christandl, K. Chang, A. Ekert, L. C. Kwek, and C. H. Oh, Phys. Rev. A 67, 012310 (2003); T. Durt, D. Kaszlikowski, J. L. Chen, and L. C. Kwek, Phys. Rev. A 69, 032313 (2004); C. Brukner, M. Żukowski, and A. Zeilinger, Phys. Rev. Lett. 89, 197901 (2002); C. Brukner, M. Żukowski, J.-W. Pan, and A. Zeilinger, Phys. Rev. Lett. 92, 127901 (2004).

d	2	3	4	5	6	7	8	9	10	20	30	40	50	60
$I_d(\|\Psi\rangle_{\rm eig})$	2.82843	2.9149	2.9727	3.0157	3.0497	3.0777	3.1013	3.1217	3.1396	3.2492	3.3068	3.3449	3.3728	3.3946
$I_d(\|\Psi\rangle_{\rm app})$	2.82843	2.90909	2.96466	3.00689	3.04078	3.06895	3.09296	3.10639	3.13219	3.24551	3.30496	3.34389	3.37222	3.39417
$I_d(\|\Psi\rangle_{\rm mes})$	2.82843	2.87293	2.89624	2.91054	2.92020	2.92716	2.93241	2.93651	2.93980	2.95472	2.95974	2.96225	2.96376	2.96477
d	70	80	90	100	150	200	250	300	350	400	450	500	550	600
$I_d(\|\Psi\rangle_{\rm eig})$	3.4124	3.4273	3.4400	3.4511	3.4914	3.5178	3.5370	3.552	3.5641	3.5743	3.583	3.5906	3.5973	3.6033
$I_d(\|\Psi\rangle_{\rm app})$	3.41192	3.4267	3.4393	3.45022	3.48933	3.51446	3.53256	3.54651	3.55775	3.5671	3.57505	3.58194	3.588	3.59339
$I_d(\|\Psi\rangle_{\rm mes})$	2.96549	2.96603	2.96645	2.96678	2.96779	2.96830	2.96860	2.96880	2.96895	2.96906	2.96914	2.96921	2.96926	2.96931
d	650	700	750	800	850	900	950	1000	1500	2000	2500	3000	3500	4000
$I_d(\|\Psi\rangle_{\rm eig})$	3.6087	3.6136	3.6181	3.6222	3.6260	3.6296	3.6329	3.6360	3.6594	3.6747	3.6859	3.6946	3.7017	3.7077
$I_d(\|\Psi\rangle_{\rm app})$	3.59824	3.60263	3.60664	3.61032	3.61372	3.61687	3.61981	3.62256	3.64299	3.65623	3.66584	3.67330	3.67936	3.68442
$I_d(\|\Psi\rangle_{\rm mes})$	2.96935	2.96938	2.96941	2.96944	2.96946	2.96948	2.96950	2.96951	2.96961	2.96966	2.96969	2.96971	2.96973	2.96974
d	5000	6000	7000	8000	50000	70000	80000	90000	100000	200000	300000	400000	500000	600000
$I_d(\|\Psi\rangle_{\rm eig})$	3.7174	3.7250	3.7311	3.7362										
$I_d(\|\Psi\rangle_{\rm app})$	3.69253	3.69884	3.70398	3.70829	3.75659	3.76372	3.76644	3.76878	3.77083	3.78345	3.79019	3.79472	3.79810	3.80080
$I_d(\|\Psi\rangle_{\rm mes})$	2.96975	2.96976	2.96977	2.96978	2.96981	2.96981	2.96981	2.96981	2.96981	2.96981	2.96981	2.96981	2.96981	2.96981

TABLE I: [Optional Table (for referee)] Bell expressions $I_d(|\Psi\rangle_{\rm eig})$, $I_d(|\Psi\rangle_{\rm app})$ and $I_d(|\Psi\rangle_{\rm mes})$.